Contents Volume 12 (1987)

Vertical distribution of radiolaria in the eastern equatorial Atlantic: analysis of a multiple series of closely-spaced plankton tows
B.A. Dworetzky and J.J. Morley (Palisades, NY, U.S.A.) 1
The distribution of benthic foraminifera in the Adriatic Sea
F.J. Jorissen (Utrecht, The Netherlands) 21
Diatom flux in McMurdo Sound, Antarctica
A. Leventer and R.B. Dunbar (Houston, TX, U.S.A.) 49
Polycystine radiolaria of the California Current region: seasonal and geographic patterns
D. Boltovskoy (Buenos Aires) and W.R. Riedel (La Jolla, CA, U.S.A.) 65

Book Reviews 105
Temporal changes in foraminiferal faunas and impact of pollution in Sandebukta, Oslo Fjord
J. Nagy and E. Alve (Oslo, Norway) 109
Foraminiferal stratigraphy and environment of Late Weichselian deposits on the continental shelf off Troms, northern Norway
M. Hald and T.O. Vorren (Tromsø, Norway) 129
Benthic foraminiferal assemblage response to the onset of northern hemisphere glaciation: paleoenvironmental changes and species trends in the northeast Atlantic
P. Loubere and G. Banonis (DeKalb, IL, U.S.A.) 161
Multivariate morphometric differentiation of chronospecies in the late Neogene planktonic foraminiferal lineage *Globoconella*
Kuo-Yen Wei (Narragansett, RI, U.S.A.) 183
Microfossil distributions in modern Amundsen Sea sediments
D.E. Kellogg and T.B. Kellogg (Orono, ME, U.S.A.) 203
Sedimentological and micropaleontological detection of turbidite muds in hemipelagic sequences: an example from the late Pleistocene levee of Monterey Fan, central California continental margin
C.A. Brunner (Berkeley, CA, U.S.A.) and M.T. Ledbetter (Moss Landing, C.A., U.S.A.) 223
Local and global effects on the Early Pliocene Mediterranean stable isotope records
L. Gudjonsson (Utrecht, The Netherlands) 241
Late Neogene paleoceanography of the eastern Caribbean, the Gulf of Mexico, and the eastern Equatorial Pacific
S. Gartner (College Station, TX, U.S.A.), J. Chow (Maoli, Taiwan) and R.J. Stanton Jr. (College Station, TX, U.S.A.) 255
Morphological and physiological responses of *Globigerinoides sacculifer* (Brady) under varying laboratory conditions
C. Hemleben (Tübingen, F.R.G.), M. Spindler (Bremerhaven, F.R.G.), I. Breitinger (Tübingen, F.R.G.) 305
Pleistocene paleoceanographic correlations. Northern Greenland Sea to central Arctic Ocean
R.S. Marquard and D.L. Clark (Madison, WI, U.S.A.) 325
Shape changes in *Globorotalia truncatulinoides* as a function of ontogeny and paleobiogeography in the Southern Ocean
R. Baxter Phar Jr. and D.F. Williams (Columbia, SC, U.S.A.) 343
Ontogeny of extant spinose planktonic foraminifera (Globigerinidae): a concept exemplified by *Globigerinoides sacculifer* (Brady) and *G. ruber* (d'Orbigny)
G.-J. A. Brummer (Amsterdam, The Netherlands), C. Hemleben (Tübingen, F.R.G.) and M. Spindler (Bremerhaven, F.R.G.) 357
Short Communication
Calpionellids and the late Jurassic and early Cretaceous stratigraphy of Kuwait and the Gulf region
I.A.Al-Rifaiy (Kuwait, Kuwait) and D. LeMone (El Paso, TX, U.S.A.) 383
Book Review 389